Form PTO-144 Patent and Trademark Office ent of Commerce

INFORMATION DISCLOSURE STATEMENT

BY APPLICANT

(Use several sheets if necessary)

Atty. Docket No. 32759	Page 1 of 1					
1 -	Serial No. 09/478,737					
Applicants	Examiner					
Wilson et. al.	Murphy					
Filing Date	Group Art Unit					
January 6, 2000	1646					

OTHER DOCUMENTS (including Author, Title Date, Pertinent Pages, Etc.) Peral et al., 1995, "Splicing mutations of the polycystic kidney disease 1 (PKD1) gene ir by intronic deletion," Human Molec. Genet. 4:569-574.	*Exam. Init.	No.			Doc	ument	No.		1	Date	Name	Class	Subclass	Filing Date if Appropriate		
OTHER DOCUMENTS (including Author, Title Date, Pertinent Pages, Etc.) Peral et al., 1995, "Splicing mutations of the polycystic kidney disease 1 (PKD1) gene in by intronic deletion," Human Molec. Genet. 4:569-574. The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease a 14kb transcript and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of PKD1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907																
Document No. OTHER DOCUMENTS (including Author, Title Date, Pertinent Pages, Etc.) Peral et al., 1995, "Splicing mutations of the polycystic kidney disease 1 (PKD1) gene in by intronic deletion," Human Molec. Genet. 4:569-574. The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease and encodes a 14kb transcript and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of PKD1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exsigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907																
OTHER DOCUMENTS (including Author, Title Date, Pertinent Pages, Etc.) Peral et al., 1995, "Splicing mutations of the polycystic kidney disease 1 (PKD1) gene in by intronic deletion," Human Molec. Genet. 4:569-574. The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease encodes a 14kb transcript and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of PKD1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907																
Document No. OTHER DOCUMENTS (including Author, Title Date, Pertinent Pages, Etc.) Peral et al., 1995, "Splicing mutations of the polycystic kidney disease 1 (PKD1) gene in by intronic deletion," Human Molec. Genet. 4:569-574. The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of PKD1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907																
OTHER DOCUMENTS (including Author, Title Date, Pertinent Pages, Etc.) Peral et al., 1995, "Splicing mutations of the polycystic kidney disease 1 (PKD1) gene in by intronic deletion," Human Molec. Genet. 4:569-574. The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease encodes a 14kb transcript and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of PKD1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907		<u> </u>			·				FO	REIGN PATI	ENT DOCUMENT			.	- '	,
OTHER DOCUMENTS (including Author, Title Date, Pertinent Pages, Etc.) Peral et al., 1995, "Splicing mutations of the polycystic kidney disease 1 (PKD1) gene in by intronic deletion," Human Molec. Genet. 4:569-574. The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease encodes a 14kb transcript and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of PKD1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Donga et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907					ocume	ent No	-						Class	SubClass	<u>Trans</u>	lator No
Peral et al., 1995, "Splicing mutations of the polycystic kidney disease 1 (PKD1) gene in by intronic deletion," Human Molec. Genet. 4:569-574. The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease gene encodes a 14kb transcript and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of PKD1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exsigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907				Γ	l	110.							Class	Buociass	165	No
Peral et al., 1995, "Splicing mutations of the polycystic kidney disease 1 (PKD1) gene in by intronic deletion," Human Molec. Genet. 4:569-574. The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease gene encodes a 14kb transcript and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of PKD1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907						-	<u> </u>									
Peral et al., 1995, "Splicing mutations of the polycystic kidney disease 1 (PKD1) gene in by intronic deletion," Human Molec. Genet. 4:569-574. The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease gene encodes a 14kb transcript and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of PKD1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exsigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907					 							<u> </u>	1	 		
Peral et al., 1995, "Splicing mutations of the polycystic kidney disease 1 (PKD1) gene in by intronic deletion," Human Molec. Genet. 4:569-574. The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease gene encodes a 14kb transcript and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of PKD1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exsigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907		L	L <u></u>	l	l	<u> </u>	<u> </u>				<u> </u>		<u> </u>	1		
by intronic deletion," Human Molec. Genet. 4:569-574. The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease encodes a 14kb transcript and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of PKD1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exsigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907	7	A		Pera										(D1) gene	induce	
gene encodes a 14kb transcript and lies within a duplicated region on chromosome 16," (77:881-894. Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of <i>PKD</i> 1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large <i>TSC2</i> and <i>PKD1</i> gene deletion is associated with renal and exsigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907			·									icy disca	se i (Fr	(1) gene	muuct	zu
Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of <i>PKD</i> 1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large <i>TSC2</i> and <i>PKD1</i> gene deletion is associated with renal and exsigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907	7	В		The European Polycystic Kidney Disease Consortium, 1994, "The polycystic kidney disease 1"												
Watnick et al., 1997, "An unusual pattern of mutation in the duplicated portion of <i>PKD</i> 1 revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large <i>TSC2</i> and <i>PKD1</i> gene deletion is associated with renal and exsigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907	1			gene 77:8	ence 81-8	odes 94	a 14	kb tr	ansc	ript and lies	within a duplicated	region on	chrom	osome 16,	" Cell	
revealed by use of a novel strategy for mutation detection," Human Molecular Genetics (1481. Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exsigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907	/	С					, 199	7, "	An u	nusual patter	n of mutation in the	duplicat	ed porti	on of PKI	01 is	.::
Longa et al., 1997, "A large TSC2 and PKD1 gene deletion is associated with renal and exsigns of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907	9			reve	aled	by u	se of	a no	vel s	strategy for n	nutation detection,"	Human N	/lolecul	ar Genetic	s <u>6</u> :147	73-
signs of autosomal dominant polycystic kidney disease," Nephrology Dialysis Transplan 12:1900-1907	₩	0				ol 1	007	66 A	lorge	TCC2 and D	OVD I cono deletion	ia assaist		11	4	
<u>// 12</u> :1900-1907	8			signs of autosomal dominant polycystic kidney disease." Nephrology Dialysis Transplantation												ena. n
Thomas et al., 1999, "Identification of mutations in the repeated part of the autosomal do				<u>12:1</u>	900-	1907	<i>'</i>									
polycystic kidney disease type 1 gene, PKD1, by long-range PCR," Am. J. Human Gene	α	E		Tho	nas (et al.	, 199	9, "I	dent	ification of n	nutations in the repe	ated part	of the a	utosomal	domin	ant

NY02:513232.1

4.29-05